



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Sunlight exposure increased Covid-19 recovery rates: A study in the central pandemic area of Indonesia



Al Asyary*, Meita Veruswati


Department of Environmental Health, Faculty of Public Health, Universitas Indonesia, Depok, Indonesia

Department of Public Health Science, Faculty of Health Sciences, Universitas Muhammadiyah Prof. Dr. HAMKA (UHAMKA), Jakarta, Indonesia

HIGHLIGHTS

- Covid-19 is a type of influenza virus that suppresses the body's immune system.
- Sunlight triggered the vitamin D that increased body immune.
- Sunlight correlated significantly to Covid-19 patients recovery in Jakarta, Indonesia

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 16 April 2020

Received in revised form 21 April 2020

Accepted 24 April 2020

Available online 27 April 2020

Editor: Wei Huang

Keywords:

Coronavirus

Sunlight

Recovery

Indonesia

ABSTRACT

This study aims to present the correlation between sunlight exposure and Covid-19 statuses in Jakarta, Indonesia. The secondary data analysis was derived from surveillance data for Covid-19 from government authorities, including the Ministry of Health, the Meteorological, Climatological, and Geophysical Agency, and the local government of Jakarta. Three statuses related to Covid-19 were examined in the study: incidence, death, and recovered. Meanwhile, sunlight exposure was presented as daily duration of it. Only the number of recovered patients correlated significantly with sunlight exposure (p -value = .025; r = 0.350). This study's findings showed that sunlight exposure was associated with recovery from Covid-19.

© 2020 Elsevier B.V. All rights reserved.

* Corresponding author at: Department of Environmental Health, Faculty of Public Health, Universitas Indonesia, C Building 2nd Floor, Kampus FKM-UI, Depok 16424, Indonesia.

E-mail address: al.asyary@ui.ac.id (A. Asyary).

1. Introduction

Coronavirus disease 2019 (Covid-19) is not only rapidly spreading but has become a global pandemic that affects many, including

developing countries with limited resources (Singhal, 2020). In fact, even the currently affected developed countries are overwhelmed in their fight against Covid-19 (Bénassy-Quééré et al., 2020; Heymann and Shindo, 2020; Soglietto et al., 2020). It is considered that most developing countries will face more challenges than developed countries in curbing the spread of Covid-19 in their regions, causing them to potentially evolve into the new epicenter of the pandemic, and Indonesia is not exempted (Hopman et al., 2020).

Covid-19 is caused by a type of influenza virus that suppresses the body's immune system so that the viruses can grow in the respiratory tissues and organs (Shi et al., 2020; Xu et al., 2020). In this situation, individuals who are elderly or with prior comorbidities are more vulnerable to the severe effects of Covid-19 due to inadequate immune systems (Garnier-Crussard et al., 2020; Le Couteur et al., 2020; Morley and Vellas, 2020). Besides that, inadequate healthcare management, such as failure to provide early diagnosis and prompt treatment, would decrease the rate of recovery among Covid-19 patients (Greenhalgh et al., 2020; Hopman et al., 2020).

Indonesia is one of the developing countries with reported Covid-19 cases (Indonesian National Task Team Force for Coronavirus 2019 (Covid-19), 2020; Organization, 2020a). International experts have expressed their doubt to the Indonesian government regarding the capability of the nation's healthcare system to control this pandemic since the authority reported the first multiple positive cases of Covid-19 on 2nd March 2020 (Djalante et al., 2020; Wenham et al., 2020). Moreover, Indonesia has one of the highest proportions of death related to Covid-19 cases among countries in the world, with the epicenter of the disease being its capital, Jakarta (Organization, 2020a).

The confirmed positive cases of Covid-19 in Indonesia are divided into three levels of severity: 1) high, 2) moderate, and 3) low (Indonesian Ministry of Health, 2020a). Patients whose conditions are of high or moderate severity are given intensive medical care in healthcare facilities. Meanwhile, low-severity cases of Covid-19 are managed the same way as the rest of the general population, meaning they should stay home and practice social isolation and physical distancing (Indonesian Ministry of Health, 2020a). At the same time, the government has strongly promoted the Clean and Healthy Living Behavior (PHBS) and People's Healthy Lifestyle Movement (Germas) programs to encourage the people to include routine outdoor/indoor exercise to decrease the spread of Covid-19 (Indonesian Ministry of Health, 2020b).

The World Health Organization (WHO) reported that there is still no evidence that sunlight can terminate Covid-19 (World Health Organization, 2020). Moreover, too much of sunlight may cause harm through its ultraviolet, infrared, etc. (Karapiperis et al., 2020). However, sunlight contributes to the healing of multiple health condition including respiratory diseases such as influenza and severe acute respiratory syndrome (SARS) (Geier et al., 2018; Miller, 2018). Weather factors have shown persistent correlation to the emergence and re-emergence of infectious diseases, including Covid-19 (Tosepu et al., 2020). Thus, evidence for the effect of sunlight on Covid-19 is needed as the correlation between the two is still rarely assessed.

2. Methods

2.1. Study area

Jakarta is the capital city and business center of Indonesia. It is geographically located within the range of 5°19'12" south latitude to 6°23'54" south latitude and 106°22'42" east longitude to 106°58'18" east longitude, with an average altitude of ± 7 m above the sea surface. Jakarta, which has tropical climates, has two seasons: rainy season that happens annually from October to March and dry season that takes place from April to September annually.

There are 132 referral hospitals available to treat Covid-19 patients of high and moderate severity in Indonesia. Eight of these, plus an

emergency hospital, are in Jakarta. They are Dr. Sulianti Saroso Central General Hospital of Infectious Disease, Persahabatan Central General Hospital of Pulmonary, Gatot Subroto Army Central Hospital, Fatmawati Central General Hospital, Tarakan Regional General Hospital, Pasar Rebo Regional General Hospital, Bhayangkara Said Soekanto Police Central Hospital, Mintoharjo Navy Central Hospital, and National Emergency Hospital of Kemayoran. These health facilities were built according to the National Hospital Accreditation Commission (KARS), including the availability of daylight exposure in each treatment unit (National Hospital Accreditation Commission, 2012).

2.2. Data collection

The set of relevant data, ranging from 2nd March to 10th April 2020, was obtained from the Indonesian government authorities. The daily confirmed cases of Covid-19 incidence, death, and recovery were derived from the Jakarta local government. This data was also made available by the Indonesian Ministry of Health as well as the Indonesian National Task Force Team for Covid-19. Meanwhile, the dataset of daily sunlight in Jakarta was obtained from the Meteorological, Climatological, and Geophysical Agency of Indonesia.

2.3. Data analysis

Spearman's correlation was performed to analyze the coefficient correlation value and determine whether significance existed. This statistical analysis was used to determine correlation between the existing variables. These variables were number of Covid-19 positive cases, deaths, and patients who recovered, and sunlight exposure.

3. Results

According to Fig. 1(a), the incidence of Covid-19 in Jakarta had been increasing. Since the first multiple cases of Covid-19 were reported on 3rd March 2020, Jakarta had seen an average of 45.25 new confirmed Covid-19 cases daily. The daily average number of recovered patients was 1.50, which was lower than the daily average deaths at 3.80 cases. Therefore, the death proportion of Covid-19 cases in Jakarta was 8.4% higher than the recovered proportion at 3.3%.

Fig. 1(b) shows that the duration of sunlight exposure for Covid-19 patients in Jakarta fluctuated daily. The shortest duration of daily sunlight exposure was 0 min, while the longest duration was 480 min in Jakarta. The average duration of daily sunlight exposure was 217.95 min or 3.6 h.

Table 1 shows that sunlight exposure did not correlate significantly with both incidence and death cases of Covid-19 patients. Sunlight correlated significantly with cases of recovery from Covid-19.

4. Discussion

In this study, we found that higher duration of sunlight exposure was related to more cases of recovery from Covid-19 among patients. This correlation is in line with previous evidence that sunlight does not terminate the Covid-19 virus so it cannot prevent the infection. Conversely, sunlight can maintain the health condition of Covid-19 patients so they have opportunity to recover from the disease. Sunlight boosts the immune system, which slows down the development of influenza and SARS agents in the human body (Cannell et al., 2006; Miller, 2018). In this case, Covid-19 patients who experienced sunlight exposure when they were receiving care either in hospitals or home settings were more likely to recover from the disease.

Sunlight triggers vitamin D production, which functions to increase the immune system (Slusky and Zechhauser, 2018). Poor sunlight exposure activates influenza (Sagripanti and Lytle, 2007). Previous studies have shown that sunlight exposure significantly contributed to the




Fig. 1. (a) Covid-19 status, (b) sunlight exposure duration (minutes).

healing of most respiratory conditions including tuberculosis and lung disease (Aloia and Li-Ng, 2007; Asyary et al., 2017).

Being in a region with a tropical climate, Jakarta can take advantage of the presence of abundant sunlight in every season. The health benefits of exposure to sunlight have been established (Singhal, 2020). Access to much-needed sunlight can enhance mood as well as release endorphins that will increase immunity against diseases such as Covid-19 (Sternberg and Engineer, 2020).

As it sits on the equator, the Indonesian territory, including Jakarta, is known as a tropical region that experiences two seasons: dry and rainy. However, even with its rainfall during the rainy season, Jakarta is still quite dry. The average duration of maximum sunlight is 3 to 3.5 h per day in Jakarta, which amounts to 4.8 kWh-hour per square meter per day ($\text{kWh}/\text{m}^2/\text{day}$), or 10 times the potential of European sunlight (Indonesia National Energy Board, 2013).

Most countries around the world, including Indonesia, are welcoming the dry or summer season, which takes place from April to September and can be potentially leveraged by each country to fight Covid-19. This potential approach can be accompanied by the formal treatment/therapy according to Covid-19 clinical guidelines (Organization, 2020b). WHO has only underlined the benefit of sunlight for clothes management (Organization, 2020c), so this finding will assist medical care particularly in low-resource health settings or places with inadequate Covid-19-mitigating response, including Jakarta (Hopman et al., 2020; Le Couteur et al., 2020).

Despite this study's significant finding about the relationship between sunlight and Covid-19 recovery, this research has limitations. The Covid-19 recovery factors could be affected by several others variables such as prompt treatment/therapy, PHBS and Germas, and

physical capabilities. It is also essential to explore the type and characteristics of sunlight exposure in detail. Additionally, the available data provided by the authorities did not record the exposure of every Covid-19 patient to sunlight.

5. Conclusion

This study shows that sunlight exposure correlated significantly with recovery from Covid-19 among patients in Jakarta of Indonesia. However, sunlight exposure did not correlate significantly with the occurrence of and death from Covid-19. This preliminary study needs to be developed further as sunlight presents genuine potential in accelerating recovery from Covid-19 not only in Jakarta but also other countries, especially those with high death rates due to this pandemic.

CRedit authorship contribution statement

AI Asyary: Conceptualization, Methodology, Data curation, Software, Writing - original draft. **Meita Veruswati:** Visualization, Validation, Writing - review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thanks to Indonesian government authorities for Covid-19, as well as to Directorate of Research and Community Engagement of Universitas Indonesia (DRPM-UI), that made the data available and supported to this study.

References

- Aloia, J.F., Li-Ng, M., 2007. RE: epidemic influenza and vitamin D epidemiology and infection October 2007, Vol. 135, no. 7, pp. 1095–1098. *Epidemiol. Infect.* 135, 1095–1098.
- Asyary, A., Eryando, T., Purwentyastuti, P., Junadi, P., Clark, C., Teijlingen, E. Van, 2017. Level of exposure of childhood tuberculosis with adult pulmonary tuberculosis

Table 1

Spearman's correlation coefficient between Covid-19 and sunlight exposure.

Confirmed Covid-19 status	Spearman's correlation coefficient
Incidence	0.306
Death	0.284
Recovery	0.350*

* Correlation is significant at the 0.05 level (2-tailed).

- household contacts. *Kesmas Natl. Public Heal. J.* 12, 1–6. <https://doi.org/10.21109/kesmas.v12i1.1469>.
- Bénassy-Quéré, A., Marimon, R., Pisani-Ferry, J., Reichlin, L., Schoenmaker, D., Weder, B., 2020. 13 COVID-19: Europe needs a catastrophe relief plan. *Mitigating COVID econ. Cris. Act Fast Do Whatever* 121.
- Cannell, J.J., Vieth, R., Umhau, J.C., Holick, M.F., Grant, W.B., Madronich, S., Garland, C.F., Giovannucci, E., 2006. Epidemic influenza and vitamin D. *Epidemiol. Infect.* 134, 1129–1140.
- Djalante, R., Lassa, J., Setiamarga, D., Mahfud, C., Sudjatma, A., Indrawan, M., Haryanto, B., Sinapoy, M.S., Rafflesia, I., Djalante, S., 2020. Review and analysis of current responses to COVID-19 in Indonesia: period of January to March 2020. *Prog. Disaster Sci.* 100091.
- Garnier-Crussard, A., Forestier, E., Gilbert, T., Krolak-Salmon, P., 2020. Novel Coronavirus (COVID-19) epidemic: what are the risks for older patients? *J. Am. Geriatr. Soc.* 1–2.
- Geier, D.A., Kern, J.K., Geier, M.R., 2018. A longitudinal ecological study of seasonal influenza deaths in relation to climate conditions in the United States from 1999 through 2011. *Infect. Ecol. Epidemiol.* 8, 1474708.
- Greenhalgh, T., Wherton, J., Shaw, S., Morrison, C., 2020. Video Consultations for Covid-19. Heymann, D.L., Shindo, N., 2020. COVID-19: what is next for public health? *Lancet* 395, 542–545.
- Hopman, J., Allegranzi, B., Mehtar, S., 2020. Managing COVID-19 in low-and middle-income countries. *JAMA* E1–E2.
- Indonesia National Energy Board, 2013. Indonesia is high potential to sunlight beneficiary [WWW Document]. News <https://www.den.go.id/index.php/dinamispage/index/396-.html>, Accessed date: 10 April 2020.
- Indonesian Ministry of Health, 2020a. National Health Care Guideline for Coronavirus 2019 (Covid-19) in Indonesia (Jakarta).
- Indonesian Ministry of Health, 2020b. Prevent Covid-19 with Germas (Jakarta).
- Indonesian National Task Team Force for Coronavirus 2019 (Covid-19), 2020. Coronavirus 2019 (Covid-19) in Indonesia (Jakarta).
- Karapiperis, C., Kouklis, P., Papastratos, S., Chasapi, A., Ouzounis, C., 2020. Assessment for the Seasonality of Covid-19 Should Focus on Ultraviolet Radiation and Not 'Warmer Days'.
- Le Couteur, D.G., Anderson, R.M., Newman, A.B., 2020. COVID-19 is a disease of older people. *J. Gerontol. A Biol. Sci. Med. Sci.* 1–5.
- Miller, B., 2018. Immune System: Your Best Defense against Viruses and Bacteria from the Common Cold to the SARS Virus. Oak Publication Sdn Bhd.
- Morley, J.E., Vellas, B., 2020. COVID-19 and older adult. *J. Nutr. Health Aging* 1.
- National Hospital Accreditation Commission, 2012. Hospital Accreditation Assessment (Jakarta).
- Organization, W.H, 2020a. Coronavirus Disease 2019 (COVID-19): Situation Report. p. 72.
- Organization, W.H, 2020b. Infection Prevention and Control Guidance for Long-Term Care Facilities in the Context of COVID-19: Interim Guidance, 21 March 2020. World Health Organization.
- Organization, W.H, 2020c. Water, Sanitation, Hygiene and Waste Management for COVID-19: Technical Brief, 03 March 2020. World Health Organization.
- Saglietto, A., D'Ascenzo, F., Zoccai, G.B., De Ferrari, G.M., 2020. COVID-19 in Europe: the Italian lesson. *Lancet* 395, 1110–1111.
- Sagripanti, J., Lytle, C.D., 2007. Inactivation of influenza virus by solar radiation. *Photochem. Photobiol.* 83, 1278–1282.
- Shi, Y., Wang, Y., Shao, C., Huang, J., Gan, J., Huang, X., Bucci, E., Piacentini, M., Ippolito, G., Melino, G., 2020. COVID-19 Infection: The Perspectives on Immune Responses.
- Singhal, T., 2020. A review of Coronavirus Disease-2019 (COVID-19). *Indian J. Pediatr.* 87, 281–286. <https://doi.org/10.1007/s12098-020-03263-6>.
- Slusky, D., Zeckhauser, R.J., 2018. Sunlight and Protection against Influenza. National Bureau of Economic Research.
- Sternberg, E., Engineer, A., 2020. Is COVID-19 making you stay at home or “shelter-in-place”? Turn your home into a healing space! [WWW document]. Andrew Weil Cent. Integr. Med. URL. https://integrativemedicine.arizona.edu/COVID19/sheltering_in_place.html, Accessed date: 10 April 2020.
- Tosepu, R., Gunawan, J., Effendy, D.S., Lestari, H., Bahar, H., Asfian, P., 2020. Correlation between weather and Covid-19 pandemic in Jakarta. Indonesia. *Sci. Total Environ.* 725, 1–4 138436.
- Wenham, C., Smith, J., Morgan, R., 2020. COVID-19: the gendered impacts of the outbreak. *Lancet* 395, 846–848.
- World Health Organization, 2020. Fact or Fiction [WWW Document]. Nov. Coronavirus 2019. URL. <https://www.who.int/southeastasia/outbreaks-and-emergencies/novel-coronavirus-2019/fact-or-fiction>, Accessed date: 11 April 2020.
- Xu, B., Fan, C., Wang, A., Zou, Y., Yu, Y., He, C., Xia, W., Zhang, J., Miao, Q., 2020. Suppressed T Cell-Mediated Immunity in Patients with COVID-19: A Clinical Retrospective Study in Wuhan, China (China (3/16/2020)).